
NANCY GRACE ROMAN





the Nancy Grace Roman Telescope


Rachael L. Beaton Roman Science Operations Center Space Telescope Science Institute





Roman Observatory

Telescope: 2.4 m aperture




<u> Wide Field Imager (WFI)</u>

- Visible / Near IR (0.48 2.3 mm)
- Field of view 0.281 deg²
- 18 4K x 4K detectors
- Grim (500), Prism (100)

<u>Coronagraph</u>

- Visible bandpass
- Contrast 10⁻⁸ 10⁻¹⁰





Mission Duration: 5 years, 10 years goal



Example Implementation of Roman Core Surveys

S 15 min cadence over few deg² Galactic bulge

Galactic Bulge Time Domain Survey


Galactic Plane

~1000 deg² in three bands (~JHK)

Galactic Plane Survey

Wide area multiband survey with spectroscopy

High Latitude Wide Area Survey



ROMAN SPACE TELESCOPE

Core Surveys



~2000 deg²

High Latitude Time Domain Survey

Tiered multiband


~10s of deg² with

time domain

spectroscopy

Roman Space Telescope's larger view and fast survey speeds will unveil the evolving universe in ways that have never been possible before.

Roman Observational Program



Astrophysics with Wide Field IR Surveys

General Astrophysics Surveys at least 25% of observing time – includes Galactic Plane Survey

- Include Galactic Plane Survey (Early Definition Community Call)
- Primarily selected via a peer-review process

Archival Investigations

- All data will be public immediately
- Anticipated to be main component of community involvement

Also
Coronagraph
Technology
Demonstration
Observations

Community definition of the Core Survey and Community Contributed Survey


Maximize the overall science return of Roman WFI IR surveys while meeting Mission Requirements focused on cosmology and exoplanets

The existing survey strategies served their primary function in showing the mission can meet its requirements.

The actual surveys to be implemented are being defined by the science community.

There will be no "survey teams" selected to define or implement the surveys.


Community definition of the Core Survey and Community Contributed Survey




Evaluate initial community input; solicit additional, targeted community input through a variety of channels; evaluate survey options against science metrics; produce recommendations for survey implementations with options for enhancements/descopes


Survey Science Helps Your Science

- Survey Speed (1000x's faster)
- Calibration Levels
- Rapid slew & settle times
- No Earth Occultation
- No SAA


















Akeson et al. 2019


Stringent Mission Calibration Requirements






Paβ in simulated stellar spectrum (Ryan+ 2019); 0.027% error

0.1% Photometric stability (for microlensing; maps to




Simulated microlensing event (S. Carey)

0.05% PSF shape (cosmic shear measurements)




Expected shear power spectrum (SDT report, Spergel+ 2015)


0.5% Absolute color calibration (for SNe; maps to luminosity distance vs. redshift)





Requirements significantly exceed what we've achieved on Hubble and the requirements for Webb





Planning Observations – Simulation tools & utilities















Planning Observations – Roman APT







Big Data



BIG DATA




Terabytes

Hubble's data archive 30 years (1990–2020)

20,000

Terabytes

Roman's data archive 5 year primary mission (projected)



Roman Science Platform Vision



The Roman Science Platform will enable the community to perform transformational science on big data by providing a rich computing environment that will allow broad, low-barrier access to data, computing, and software resources.

- Enable collaboration
- Support the community
- Build software and tools that bring people to the platform

Roman Science Platform Vision




- Cloud-compute environments Jupyterlab
- Brings users to the data easy and efficient access, decreased need for downloads
- Cloud-based computing resource

Resources for exploratory work will be available with no need to write

grant proposal

Offers flexible and scalable architecture

 Includes pre-installed software and tools for calibration, analysis, visualization, and training





Roman Documentation







Welcome to the Nancy Grace Roman Space Telescope Help Desk



SOC Roman Helpdesk

Quick Start APT for Roman





Simulation Tools Handbook

Getting involved with Roman

Roman Working Groups



Calibrations, Simulations, Software, Survey Definition

Roman ST Advisory Committee





Roman Science Interest Group

Roman conference "Challenging Theory with Roman: From Planet Formation to Cosmology", Caltech campus + online, July 9-12, 2024





Nancy Grace Roman


NASA's First Chief of Astronomy

1925-2018



Roman's Instrument Capabilities

Roman Space Telescope Imaging Capabilities													
Telescope Aperture (2.4 meter)		(4	Field of View (45'x23'; 0.28 sq deg)		eg)	Pixel Scale (0.11 arcsec)				Wavelength Range (0.5-2.3 μm)			
Filters	F0	62	F087	F087 F106		F129	F	158 F18		4	F213		W146
Wavelength (μ	m) 0.48-	0.76	6 0.76-0.98 0.93		.19 1	.13-1.45	1.38-1.77		1.68-2.00		1.95-2.3	30	0.93-2.00
Sensitivity (5σ AB mag in 1	hr) 28	.5	28.2	28.1	28.0		2	28.0 27.		7.5 26.2			28.3
Roman Space Telescope Spectroscopic Capabilities													
		Fi	Field of View (sq deg)		Wavel	elength (µm)		Resolution			Sensitivity (AB mag) (10σ per pixel in 1hr)		
Grism	Grism 0.28 sq deg			3	1.00-1.93		461			20.5 at 1.5 μm			
Prism		C).28 sq deg	eg 0		.75-1.80		80-	80-180		23.5 at 1.5 μm		
Roman Space Telescope Coronagraphic Capabilities													
	Waveleng (µm)	jth	Inner Working Angle (arcsec)		Outer Working Ang (arcsec)		gle I	Detection Limit*			pectral solution		
Imaging	0.5-0.8		0.15 (exoplanets) 0.48 (disks)			0.66 (exoplanets) 1.46 (disks)) '	10 ⁻⁹ contrast (after post- processing)		47-75		
Spectroscopy	0.675-0.7	85						,			47-75		



Galactic Plane Survey

Roman's high angular resolution, sensitivity, NIR wavelengths, and survey speed will minimize confusion and extinction in the plane of the Galaxy, enabling surveys that can

- Resolve previously unresolved stellar populations, e.g., stellar clusters in star-forming regions, globular clusters
- Place new constraints on Galactic structure, by surveying red clump stars and YSOs over a greater volume of the disk
- Measure the stellar Initial Mass Function to lower mass limits; resolve low mass stars in extincted regions
- Observe more "background" sources for the construction of 3D dust maps
- Obtain a complete census of evolved massive stars in the disk: measure physical properties, constrain their binary fraction, and enable multimessenger stellar astronomy

