

# ExoPAG SAG 23

## The Impact of Exo-Zodiacal Dust on Exoplanet Direct Imaging Surveys

Co-leads


John Debes (Space Telescope Science Institute)

Isabel Rebollido (European Space Agency, European Space Astronomy Centre)

Yasuhiro Hasegawa (Jet Propulsion Laboratory, California Institute of Technology)



# What is exozodi? Why do we care?



Planetary systems are composed of planets, small bodies (e.g., asteroids, comets), and tiny dust particles


Warm and hot dust are often called as exozodiacal dust, or exozodi in short

Warm dust:

- present around habitable zones
- thermal emission around 10 micron
- becomes noise/confusion sources by scattered light in visible



Hot dust:

- present near the host star
- thermal emission around 1 micron
- introduces coronagraphic leakage in visible



Credit: S. Ertel

# What is exozodi? Why do we care?



Planetary systems are composed of planets, small bodies (e.g., asteroids, comets), and tiny dust particles

Warm and hot dust are often called as exozodiacal dust, or exozodi in short

Warm dust:


- present around habitable zones
- thermal emission around 10 micron
- becomes noise/confusion sources by scattered light in visible

Hot dust:

- present near the host star
- thermal emission around 1 micron
- introduces coronagraphic leakage in visible

Exozodi is one key parameter to affect yields of Earth-like planets by the Habitable World Observatory (HWO)!!!

# Previous findings/recommendations



various ground-based IR and sub-millimeter interferometers and a possible coronagraphic capability for WF3 (see below) on the Hubble Space Telescope (HST), to independently examine and compare the expected sensitivities, resolutions, and other important parameters. One important outcome of such a study would be the determination of how these facilities can be used together to maximize our knowledge of exozodiacal dust.

## SPECIFIC RECOMMENDATIONS

The recommendations from the Ames Exozodiacal Dust Workshop can be summarized as arguing for the establishment of a national science program to provide a solid understanding and characterization of exozodiacal and zodiacal dust emission and of the potential TPF target stars.


- There should be a program of observations that make use of available or planned facilities such as the Keck Interferometer, Large Binocular Telescope (LBT), Space Infrared Telescope Facility (SIRTF), Stratospheric Observatory for Infrared Astronomy (SOFIA), etc., to directly measure the exozodiacal emission around nearby stars. A present estimate is that a TPF interferometer resembling the present baseline designs would have degraded sensitivity to earth-like planets in systems with exozodiacal dust densities above about 10x that in our system. Systems with less dust than that become primary TPF targets. (These limits, however, will depend on the actual levels of spatial fluctuations in the exozodiacal dust clouds.)

Therefore, an attempt should be made to measure the fluctuation power spectrum in the brighter dust systems down to 0.1 AU-resolution, and then extend these results to fainter systems by modeling. Observations with the Keck Interferometer will be particularly valuable in this area.

- There should be a parallel program directed at collating what is known about all the normal stars in the solar neighborhood and at promoting the filling in and extension of that database so that target selection criteria other than exozodiacal emission strength can be understood.

- Finally, our theoretical understanding of the sources, sinks, dynamics, and evolution of our own zodiacal cloud needs to be substantially increased before we can interpret exozodiacal observations in comparison with our solar system.

These three broad programs of scientific inquiry are intrinsically interesting and should be pursued no matter what techniques and instruments are finally used to look for terrestrial planets. Support for these programs will be crucial to success in finding extrasolar terrestrial planets. At the least, support should take the form of a substantial augmentation to the existing "Origin of Solar Systems" grants program, thereby allowing a significant number of new grants, directed toward modeling of dust clouds in planetary systems.



Adopted from SAG 1 report (Roberge et al 2012)

20+ year old recommendations!

The occurrence rate of *warm* exozodi should be constrained by observations with LBTI and other facilities

All other observations available should be used to develop a reliable target selection strategy

Theoretical understanding of the origin of *Solar* zodi should be increased

# Focused areas of SAG 23

Theory of Exozodi Sources and Dust Evolution (M. Wyatt)

Pan-Chromatic Radiative Transfer of Exozodis (R. Anche)

Hot Dust (S. Ertel, W. Danchi)

A Catalog of Dusty Systems around Nearby Stars (A. Tanner, S. Ertel)

Solar System Zodi (G. Bryden, N. Turner)

Prioritization of Precursor Observational Studies of Debris Disks/Exozodis for Future Direct Imaging Missions (M. Millar-Blanchaer, W. Danchi)



Prioritization of Precursor Theoretical Studies of Debris Disks/Exozodis for Future Direct Imaging Missions (V. Faramaz, S. Dodson-Robinson)

Update and Prioritization of ExEP Gaps relevant to ExoZodis (K. Hoch)

About 40 experts of exozodi, Solar zodi, and debris disks in general contribute to these efforts

# HOST survey by LBTI at the $N$ band around 10 micron

Ertel et al. 2020





20-40 % of main-sequence stars have detectable *warm* exozodi

A positive correlation with cold debris disks is inferred


The majority of Sun-like stars have relatively low *warm* exozodi levels (best-fit median: 3 zodis)!

# Knowledge gap 1: scattered light from *warm* exozodi at visible



Current large inner working angle of visible observations prevents direct comparison between thermal emission and scattered light from *warm* exozodi


# Knowledge gap 2: the presence of *hot* exozodi and coronagraphic leakage




The presence of *hot* exozodi appears not to correlate with stellar spectral type and the presence of other dust in the system

Typical *hot* exozodi flux is around 1% of the stellar flux in the H or K bands

For some stars, hot dust resides at  $\sim \lambda/D$  for HWO, causing coronagraphic leakage, as already suggested by JWST observations




Thermal emission from *hot* Exozodi is relatively insensitive to grain size



Scattered light from *hot* Exozodi is very sensitive to grain size

# Knowledge gap 3: Poorly constrained origins of *warm* and *hot* exozodis



The theory chapter by Wyatt et al.

## *Warm* exozodi

- Inward drift of cold dust by PR drag possibly interacting with planets
- Inward scattering of exocomets by planets
- Other dynamical processes (e.g., Kozai)

## *Hot* exozodi

- Sublimation of dust
- Magnetic trapping
- Gas drag

# Key steps to fill out gap 1: better characterize *warm* exozodi

The precursor observation chapter by Millar-Blanchaer, Danchi et al.

- **Complete sensitive searches for mid-IR excesses for nearby stars at 10  $\mu\text{m}$**   
Only 38 of the 68 targets were observed by HOSTS  
A larger sample at HOSTS sensitivity could provide a constraint on the median exozodi level **three times** better than now
- **Support high contrast imaging in visible total intensity and polarized light**  
Requires  $10^{-8}$ - $10^{-10}$  contrast at  $\sim 0.1''$  inner working angle to be sensitive to 1-100x the solar system zodi

# Key steps to fill out gap 2: better characterize *hot* exozodi

The hot dust chapter by Ertel et al.

- **Observe and characterize hot exozodi using near-IR to mid-IR interferometry**  
**Dense u-v-coverage** and **spectral resolution** will provide improved constraints on the properties of hot exozodi (e.g., **dust geometry** including asymmetric structures in the spatial distribution, **spectral features**)
- **Explore luminosity function and variability using nulling beam L band observations that have contrasts of  $10^{-5}$  between 5-80 mas**  
This sensitivity will allow **ten to fifty times** more tenuous hot exozodi systems than currently possible and thus to derive a more accurate luminosity function for hot dust

# Key steps to fill out gap 1: better characterize *warm* exozodi

The precursor observation chapter by Millar-Blanchaer, Danchi et al.

- **Complete sensitive searches for mid-IR excesses for nearby stars at 10  $\mu\text{m}$**   
Only 38 of the 68 targets were observed by HOSTS  
A larger sample at HOSTS sensitivity could provide a constraint on the median exozodi level **three times** better than now
- **Support high contrast imaging in visible total intensity and polarized light**  
Requires  $10^{-8}$ - $10^{-10}$  contrast at  $\sim 0.1''$  inner working angle to be sensitive to 1-100x the solar system zodi

# Key steps to fill out gap 2: better characterize *hot* exozodi

The hot dust chapter by Ertel et al.

- **Observe and characterize hot exozodi using near-IR to mid-IR interferometry**  
**Dense u-v-coverage** and **spectral resolution** will provide improved constraints on the properties of hot exozodi (e.g., **dust geometry** including asymmetric structures in the spatial distribution, **spectral features**)
- **Explore luminosity function and variability using nulling beam L band observations that have contrasts of  $10^{-5}$  between 5-80 mas**  
This sensitivity will allow **ten to fifty times** more tenuous hot exozodi systems than currently possible and thus to derive a more accurate luminosity function for hot dust

# Key steps to fill out gap 3: improve our understanding of exozodi

The precursor theory chapter by Faramaz et al. & the pan-chromatic radiative transfer chapter by Anche et al.

- **Develop predictable models for the properties of exozodi**

Conduct a **thorough** parameter study including **various delivering mechanisms** and reveal any relationships between the properties of exozodi and planetary system architectures

- **Develop end-to-end modeling frameworks in which different telescope and instrument designs will be coupled with detailed treatments of dust scattering properties**

Construct **sophisticated grain models** based on experimental measurements and observations and infuse into **radiative transfer simulations** to better characterize observed properties (e.g., speckle shape and size) of exozodi in visible

## Completed and planned activities

- **One-day workshop** (Sept, 15, 2023 at STScI)

About 20 in person and 50 remote participants with a good mixture of career levels

Active discussions with nearly equal interests on each focused areas including talks from community members

- **Documentation**

The hot dust chapter was accepted to PASP as a review; Theory of Exozodis submitted to PASP fall 2025

Updated recommendations on the ExEP Gap list (the science gap list chapter by Hoch et al.)

HWO Science Case for ExoZodis Summer 2025; DARES white paper Spring 2025

**Final Report will be finalized soon**

# Key steps to fill out gap 3: improve our understanding of exozodi

The precursor theory chapter by Faramaz et al. & the pan-chromatic radiative transfer chapter by Anche et al.

- **Develop predictable models for the properties of exozodi**

Conduct a **thorough** parameter study including **various delivering mechanisms** and reveal any relationships between the properties of exozodi and planetary system architectures

- **Develop end-to-end modeling frameworks in which different telescope and instrument designs will be coupled with detailed treatments of dust scattering properties**

Construct **sophisticated grain models** based on experimental measurements and observations and infuse into **radiative transfer simulations** to better characterize observed properties (e.g., speckle shape and size) of exozodi in visible

## Completed and planned activities

- **One-day workshop** (Sept, 15, 2023 at STScI)

About 20 in person and 50 remote participants with a good mixture of career levels

Active discussions with nearly equal interests on each focused areas including talks from community members

- **Documentation**

The hot dust chapter was accepted to PASP as a review; Theory of Exozodis submitted to PASP fall 2025


Updated recommendations on the ExEP Gap list (the science gap list chapter by Hoch et al.)

HWO Science Case for ExoZodis Summer 2025; DARES white paper Spring 2025

**Final Report will be finalized soon**

Backup slides

# Coronagraphs like HWO will be Exozodi finding machines



## Exozodi Characterization

- Earth finding requirements are excellent for exozodi detection
- Characterization requires:
  - Broad wavelength coverage to constrain composition
  - Polarimetry to constrain grain properties
  - Spatial resolution for morphology
- Well resolved exozodis can be searched for perturbations in density due to low mass planets